

Non-cannabinoid constituents from a high potency *Cannabis sativa* variety

Mohamed M. Radwan^a, Mahmoud A. ElSohly^{a,b,*}, Desmond Slade^a, Safwat A. Ahmed^a, Lisa Wilson^c, Abir T. El-Alfy^c, Ikhlas A. Khan^{a,d}, Samir A. Ross^{a,d,*}

^a National Center for Natural Products Research, School of Pharmacy, The University of Mississippi, University, MS 38677, USA

^b Department of Pharmaceutics, School of Pharmacy, The University of Mississippi, University, MS 38677, USA

^c Department of Pharmacology, School of Pharmacy, The University of Mississippi, University, MS 38677, USA

^d Department of Pharmacognosy, School of Pharmacy, The University of Mississippi, University, MS 38677, USA

ARTICLE INFO

Article history:

Received 2 May 2008

Received in revised form 3 June 2008

Available online 4 September 2008

Keywords:

Cannabis sativa L.

Cannabaceae

High potency

Non-cannabinoid

Antimicrobial

Antileishmanial

Antimalarial

Anti-oxidant

ABSTRACT

Six new non-cannabinoid constituents were isolated from a high potency *Cannabis sativa* L. variety, namely 5-acetoxy-6-geranyl-3-*n*-pentyl-1,4-benzoquinone (**1**), 4,5-dihydroxy-2,3,6-trimethoxy-9,10-dihydrophenanthrene (**2**), 4-hydroxy-2,3,6,7-tetramethoxy-9,10-dihydrophenanthrene (**3**), 4,7-dimethoxy-1,2,5-trihydroxyphenanthrene (**4**), cannflavin C (**5**) and β -sitosterol-3-*O*- β -D-glucopyranoside-2'-*O*-palmitate (**6**). In addition, five known compounds, α -cannabispiranol (**7**), chrysoeriol (**8**), 6-prenylapigenin (**9**), cannflavin A (**10**) and β -acetyl cannabispiranol (**11**) were identified, with **8** and **9** being reported for the first time from cannabis. Some isolates displayed weak to strong antimicrobial, antileishmanial, antimalarial and anti-oxidant activities. Compounds **2–4** were inactive as analgesics.

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction

Cannabinoids are phenolic compounds possessing a C₂₁ terpenophenolic structure uniquely found in *Cannabis sativa* L. (ElSohly and Slade, 2005). Currently, 86 cannabinoids have been isolated from cannabis (Ahmed et al., 2008; ElSohly and Slade, 2005; Radwan et al., 2008). Non-cannabinoid constituents isolated from cannabis include flavonoids, spiroindans, dihydrostilbenes, dihydrophenanthrenes, sterols and alkaloids, among others (Ross and ElSohly, 1995; Turner et al., 1980). As part of our program to study the constituents of high potency cannabis and their pharmacology (Ahmed et al., 2008; Radwan et al., 2008), we herein report the isolation and structure elucidation of eleven non-cannabinoid constituents including six new (**1–6**) and five known (**7–11**) compounds as well as their antimicrobial, antileishmanial, antimalarial and anti-oxidant activities. The analgesic activities of **2–4** are also reported.



2. Results and discussions

Compound **1** was obtained as an orange amorphous powder. Its positive mode HRESIMS exhibited a pseudomolecular ion at m/z 373.2425 [M+H]⁺ corresponding to a molecular formula of C₂₃H₃₂O₄. The IR spectrum of **1** indicated the presence of an α,β -unsaturated ketone and ester carbonyl groups at ν_{\max} 1663 and 1780 cm⁻¹, respectively. The ¹H NMR spectrum of **1** displayed three olefinic methyl singlets (δ_{H} 1.56, 1.63 and 1.69), one primary methyl triplet (δ_{H} 0.87, H₃-5'), one acetoxy methyl singlet (δ_{H} 2.32), one aromatic singlet (δ_{H} 6.53, H-2) and 7 methylene resonances (δ_{H} 1.32–2.39). The ¹³C NMR, DEPT and HMQC spectra of **1** revealed 23 resonances including five methyl, seven methylene, one aromatic methine, two olefinic methine and eight quaternary carbons. The two carbonyl carbons resonating at δ_{C} 187.1 and 180.8 are characteristic for a benzoquinone skeleton (Mossa et al., 1999), while NMR spectroscopic analysis suggested an acetoxy (δ_{H} 2.32, δ_{C} 168.1, 20.5), a geranyl (Radwan et al., 2008) and an *n*-pentyl substituent (Ahmed et al., 2008; Radwan et al., 2008), indicating that **1** is a trisubstituted-1,4-benzoquinone derivative. HMBC analysis (Fig. 1) placed the geranyl substituent at C-6 due to the correlation of H₂-1" (δ_{H} 3.10) with C-6 (δ_{C} 135.6), C-1 (δ_{C} 187.1) and C-5 (δ_{C} 149.1). The three bond HMBC correlation of H-2 (δ_{H} 6.53) with C-6 (δ_{C} 135.6) and C-4 (δ_{C} 180.8), and the correlation of H₂-1' (δ_{H} 2.39) with C-4 (δ_{C} 180.8) and C-2 (δ_{C} 132.7) placed the *n*-pentyl





* Corresponding authors. Address: National Center for Natural Products Research, School of Pharmacy, The University of Mississippi, University, MS 38677, USA. Tel.: +1 662 915 1031; fax: +1 662 915 7989 (S.A. Ross), tel.: +1 662 915 5928; fax: +1 662 915 5587 (M.A. ElSohly).

E-mail addresses: melsohly@olemiss.edu (M.A. ElSohly), ssross@olemiss.edu (S.A. Ross).

moiety at C-3. The location of the acetoxy group at C-5 was determined by the four bond HMBC correlation between the acetoxy methyl (δ_{H} 2.32) and C-5 (δ_{C} 149.1). The presence of the *n*-pentyl and geranyl groups were confirmed by COSY correlations (Fig. 1), establishing **1** as 5-acetoxy-6-geranyl-3-*n*-pentyl-1,4-benzoquinone.



δ_{C} 30.6 (C-9) and 31.8 (C-10) in the HMQC spectrum], a pair of signals for an *ortho*-coupled AB spin system [δ_{H} 6.74 (1H, *d*, $J = 8.0$ Hz, H-7) and 6.79 (1H, *d*, $J = 8.0$ Hz, H-8) correlated to δ_{C} 109.2 (C-7) and 119.1 (C-8), respectively, in the HMQC spectrum], and an isolated aromatic proton [δ_{H} 6.50 (s, H-1)] suggesting that **2** is a 2,3,4,5,6-pentasubstituted 9,10-dihydrophenanthrene (Crombie



Compound **2** was isolated as a brown amorphous powder. Its molecular formula was determined as C₁₇H₁₈O₅ from the positive mode HRESIMS ion at m/z 325.1100 [M+Na]⁺ and ¹³C NMR spectrum. The ¹H and ¹³C NMR spectra of **2** (Table 1) displayed two sets of methylene protons [δ_{H} 2.64 (4H, s, H₂-9 and H₂-10) correlated to

et al., 1979; Leong et al., 1997; Stermitz et al., 1983). The ¹³C NMR (Table 1) and APT spectra indicated the presence of five oxygenated quaternary aromatic carbons. Three of these carbons have methoxy substituents as established by three sharp singlets in the ¹H NMR [δ_{H} 3.85, 3.86 and 3.89 (3H each)], while the remaining




Fig. 1. HMBC and COSY correlations of **1**.

two carbons have hydroxyl substituents as was confirmed by the presence of a characteristic absorption band at ν_{\max} 3420 cm^{-1} in the IR spectrum. The 3J HMBC correlations between H-8 (δ_{H} 6.79) and C-6 (δ_{C} 147.6), C-4b (δ_{C} 121.1) and C-9 (δ_{C} 30.6); between H-7 (δ_{H} 6.74) and C-8a (δ_{C} 132.6) and C-5 (δ_{C} 141.7) and between the methoxyl group at δ_{H} 3.85 and C-6 (δ_{C} 147.6), located this methoxyl group at C-6, as was confirmed by ROESY correlation (Fig. 2). The 3J HMBC correlation between the isolated aromatic proton at δ_{H} 6.50 and C-10 (δ_{C} 31.8) and C-4a (δ_{C} 114.9) determined its location at C-1. The ROESY correlation between the remaining two methoxyl groups at δ_{H} 3.86 and 3.89 in conjunction with the ROESY correlation between the methoxyl group at δ_{H} 3.89 and H-1 (δ_{H} 6.50) fixed their positions at C-3 and C-2, respectively, which was confirmed by HMBC correlations (Fig. 2), establishing **2** as 4,5-dihydroxy-2,3,6-trimethoxy-9,10-dihydrophenanthrene.

Table 1

^1H (400 MHz) and ^{13}C (100 MHz) NMR spectroscopic data of **2–4** (CDCl_3 , δ in ppm, J in Hz)^a

Position	2		3		4	
	δ_{H}	δ_{C}	δ_{H}	δ_{C}	δ_{H}	δ_{C}
1	6.50 s	104.4	6.39 s	103.9	–	140.2
2	–	152.0	–	150.3	–	140.2
3	–	136.4	–	134.5	6.15 s	107.6
4	–	147.2	–	146.8	–	161.5
4a	–	114.9	–	114.7	–	124.2
4b	–	121.1	–	107.8	–	117.4
5	–	141.7	8.02 s	111.5	–	156.6
6	–	147.6	–	147.2	6.93 d (2.0)	108.9
7	6.74 d (8.0)	109.2	–	147.1	–	161.0
8	6.79 d (8.0)	119.1	6.79 s	111.1	6.82 d (2.0)	102.0
8a	–	132.6	–	125.4	–	129.0
9	2.64 s	30.6	2.74 s	29.5	8.06 d (8.4)	137.7
10	2.64 s	31.8	2.74 s	30.9	8.12 d (8.4)	122.9
10a	–	136.4	–	130.1	–	132.7
OMe-2	3.89 s	56.3	3.88 s	56.2	–	–
OMe-3	3.86 s	61.0	3.89 s	61.3	–	–
OMe-4	–	–	–	–	3.96 s	57.2
OMe-6	3.85 s	56.1	3.92 s	56.0	–	–
OMe-7	–	–	3.92 s	56.1	3.90 s	55.7

^a Assignments confirmed by APT, gHMBC, gCOSY and gHMBC experiments.

Compound **3** was isolated as a pale brownish, amorphous powder. The molecular formula of **3** was determined as $\text{C}_{18}\text{H}_{20}\text{O}_5$ from the positive mode HRESIMS at m/z 339.1279 $[\text{M}+\text{Na}]^+$, a mass difference of 14 amu compared to **2**. The ^1H and ^{13}C NMR spectra of **3** (Table 1) were similar to **2** except for the replacement of one hydroxyl by a methoxyl group (δ_{H} 3.92, δ_{C} 56.1). Therefore, **3** exhibited four methoxyl groups [δ_{H} 3.88 (3H, s), 3.89 (3H, s) and 3.92 (6H, s); δ_{C} 56.0, 56.1, 56.2 and 61.3] and one hydroxyl group. Their




Fig. 2. HMBC and ROESY correlations of **2**, **5** and **6**.

locations on the dihydrophenanthrene skeleton were determined by comparison to the ^1H and ^{13}C NMR data of **2** (Table 1) and confirmed by HMBC and ROESY correlations, placing two methoxyl groups at C-2 and C-3 and the hydroxyl at C-4. The remaining methoxyl groups are therefore attached to ring B. The presence of a pair of isolated aromatic singlets [δ_{H} 6.79 (H-8) and 8.02 (H-5)] in the ^1H NMR assigned the two methoxyl groups to C-6 (δ_{C} 147.2) and C-7 (δ_{C} 147.1) (Leong et al., 1997), which was confirmed by ROESY [δ_{H} 3.92 (OMe-6 and OMe-7)/H-5 and H-8; H-8/H₂-9] and HMBC (H-5/C-7, C-4a; H-8/C-6, C-8a, C-9) correlations. Thus, the structure of **3** was established as 4-hydroxy-2,3,6,7-tetramethoxy-9,10-dihydrophenanthrene.

Compound **4** was isolated as a reddish brown powder. Its molecular formula was determined as $\text{C}_{16}\text{H}_{14}\text{O}_5$ from negative mode HRESIMS (m/z 571.1630 [2M-H] $^-$) and ^{13}C NMR data. The UV spectrum of **4** (λ_{max} 258, 282 and 303 nm) is characteristic for a phenanthrene skeleton (Leong et al., 1997). The ^1H NMR (Table 1) displayed a pair of signals for an *ortho*-coupled AB spin system [δ_{H} 8.06 (1H, *d*, $J = 8.4$ Hz, H-9), 8.12 (1H, *d*, $J = 8.4$ Hz, H-10)], two *meta*-coupled protons [δ_{H} 6.82 (1H, *d*, $J = 2.0$ Hz, H-8), 6.93 (1H, *d*, $J = 2.0$ Hz, H-6)] and an isolated aromatic singlet [δ_{H} 6.15 (1H, *s*, H-3)] indicative of a pentasubstituted phenanthrene (Réthy et al., 2006; Leong et al., 1997). The ^1H and ^{13}C NMR spectra of **4** showed two aromatic methoxyl groups [δ_{H} 3.90 (3H, *s*) and 3.96 (3H, *s*)] and five oxygenated quaternary carbons, indicating that **4** has three hydroxyl groups. The ROESY correlation of the methoxyl group at δ_{H} 3.90 with the protons at δ_{H} 6.93 (H-6) and 6.82 (H-8) established its location at C-7 (δ_{C} 161.0), which was confirmed by HMBC [δ_{H} 3.90 (OMe-7)/C-7; H-8/C-7, C-6, C-8a, C-9; H-6/C-7, C-8] correlations. The ROESY and HMBC correlations of the methoxyl group at δ_{H} 3.96 with H-3 (δ_{H} 6.15) and C-4 (δ_{C} 161.5), respectively, assigned its location at C-4. Therefore, the structure of **4** was determined as 4,7-dimethoxy-1,2,5-trihydroxyphenanthrene.

Compound **5** was obtained as a yellow amorphous powder. Its positive mode HRESIMS displayed an $[\text{M}+\text{Na}]^+$ ion at m/z 459.1766 suggesting $\text{C}_{26}\text{H}_{28}\text{O}_6$ as the molecular formula and 13 degrees of unsaturation. The IR spectrum showed absorption bands at ν_{max} 3421 and 1662 cm^{-1} due to hydroxyl and carbonyl groups, respectively, while the UV absorption maxima at λ_{max} 275 (band I) and 340 (band II) nm were indicative of a flavone skeleton (Mabry et al., 1970). The ^1H NMR spectrum of **5** established a hydrogen-bonded hydroxyl proton [δ_{H} 13.05 (*s*, HO-5)] which was confirmed by the bathochromic UV shift (+25 nm) of band II upon the addition of AlCl_3 to a methanolic solution of **5**. Bathochromic UV shifts upon the addition of NaOMe (+61 nm) and NaOAc (+5 nm) suggested hydroxylation at C-4' and C-7, respectively (Mabry et al., 1970). The ^1H NMR spectrum displayed two sharp singlets at δ_{H} 6.66 (1H, H-3) and 6.36 (1H, H-6), one methoxyl group (δ_{H} 3.98) and an ABX spin system of ring B [δ_{H} 7.01 (1H, *d*, $J = 8.0$ Hz, H-5'), 7.58 (1H, *d*, $J = 2.0$ Hz, H-2'), 7.64 (1H, *dd*, $J = 2.0, 8.0$ Hz, H-6')]. The presence of a geranyl group was deduced from the three methyl singlets at δ_{H} 1.48, 1.53 and 1.82 and two olefinic proton triplets at δ_{H} 5.02 and 5.35 in the ^1H NMR spectrum (Ahmed et al., 2008; Radwan et al., 2008). The ^{13}C NMR, DEPT-135 and HMQC spectra displayed 26 resonances including three methyl, one methoxyl, three methylene, seven methine and 12 quaternary carbons. The location of the methoxyl group was determined to be at C-3' from HMBC (OMe-3'/C-3'; H-5'/C-3'; H-2'/C-3', C-4') and ROESY (OMe-3'/H-2') correlations (Fig. 2). The carbon resonance at δ_{C} 98.9 corresponding to a proton singlet at δ_{H} 6.36 in the HMQC spectrum indicated an unsubstituted C-6 position (Agrawal, 1989). The location of the geranyl group at C-8 was confirmed by the HMBC correlation of the benzylic protons [δ_{H} 3.57 (2H, *d*, $J = 6.8$ Hz, H₂-1'')] with C-8 (δ_{C} 106.5), C-7 (δ_{C} 162.4) and C-9 (δ_{C} 155.4) (Fig. 2). The spectroscopic data of **5** are similar to those reported for cannflavin A (**10**) (Agrawal, 1989; Choi et al., 2004) except for

the location of the geranyl group at C-8 instead of C-6, establishing **5** as 8-geranyl-5,7,4'-trihydroxy-3'-methoxyflavone (cannflavin C).

Compound **6** was obtained as an optically active white amorphous powder. Its molecular formula was deduced from the positive mode HRESIMS $[\text{M}+\text{Na}]^+$ ion at m/z 837.6621 as $\text{C}_{51}\text{H}_{90}\text{O}_7$. The ^1H NMR spectrum displayed two tertiary [δ_{H} 0.75 (Me-18) and 0.99 (Me-19)], three secondary [δ_{H} 0.63 (Me-26 and Me-27) and 0.90 (Me-21)] and one primary [δ_{H} 0.63 (Me-29)] methyl groups in addition to an olefinic proton at δ_{H} 5.33 (*bs*, H-6), indicating a sitosterol skeleton (Kovganko et al., 2000; Takemoto et al., 1967). The presence of an anomeric proton [δ_{H} 4.33 (*d*, $J = 7.6$ Hz, H-1')] and carbon [δ_{C} 101.6 (C-1')] in the HMQC spectrum indicated monoglycosylation at C-3 (δ_{C} 80.1) (Ishii et al., 1977) that was confirmed by HMBC (H-3/C-1'; H-1'/C-3) and ROESY (H-1'/H-3) correlations (Fig. 2). The sugar moiety was identified as β -D-glucopyranose by acid hydrolysis of **6** and TLC comparison with authentic sugar samples. The ^1H and ^{13}C NMR spectroscopic data of **6** were similar to those reported for β -sitosterol-3-O- β -D-glucopyranoside (Chang et al., 1981) with the addition of 16 resonances characteristic for a palmitate moiety (Segre and Mannina, 1997), which was confirmed by methylation of the alkaline hydrolysis product of **6** followed by GCMS analysis. The downfield esterification shift of C-2' (+2 ppm), upfield shifts of C-1' (-4 ppm) and C-3' (-3 ppm) and the four bond HMBC correlation of H-1' (δ_{H} 4.33) and C-1'' (δ_{C} 174.6) (Fig. 2) placed the palmitate moiety at C-2' (Terui et al., 1976; Yamasaki et al., 1977). Thus, the structure of **6** was established as β -sitosteryl-3-O- β -D-glucopyranoside-2'-O-palmitate.

Compound **7** was obtained as colorless prisms. Its positive mode HRESIMS gave an $[\text{M}+\text{H}]^+$ ion at m/z 249.2393 corresponding to a molecular formula of $\text{C}_{15}\text{H}_{20}\text{O}_3$. The GCMS showed a molecular ion at m/z 248 (33%) and two characteristic ions at m/z 189 (100%) and 176 (65%) suggesting that **7** is a spiroindan derivative (Erfaly et al., 1986). The ^{13}C NMR, DEPT and HMQC spectra of **7** displayed 15 resonances including one methoxyl, six methylene, two aromatic methine, one sp^3 oxymethine and five quaternary carbons. The spectroscopic data of **7** are similar to those reported for β -cannabispiranol (Boeren et al., 1977; Shoyama and Nishioka, 1978; Radwan et al., 2008) except for the downfield shift of the oxymethine carbon (+5.6 ppm), indicating a 4'-configuration. Although **7** is a known cannabis constituent (Crombie and Crombie, 1982), this is the first report of the full NMR assignments.

The flavones **8** and **9** were isolated as yellow amorphous powders. Their molecular formulae were determined from the HRESIMS as $\text{C}_{16}\text{H}_{12}\text{O}_6$ and $\text{C}_{20}\text{H}_{18}\text{O}_5$, respectively. Their spectroscopic data (UV and NMR) were in agreement with reported values for chrysoeriol (**8**) (Toth et al., 1980; Agrawal, 1989) and 6-prenylapigenin (**9**) (Abegaz et al., 1998). This is the first report of their isolation from cannabis. The NMR spectra of **10** and **11** were identical with those of cannflavin A (Agrawal, 1989; Choi et al., 2004) and β -acetyl cannabispiranol (Shoyama and Nishioka, 1978), respectively.

The antimicrobial, antileishmanial, antimalarial and anti-oxidant activities of the isolated compounds were tested. Compound **1** displayed weak anti-MRSA (IC_{50} 15.0 $\mu\text{g}/\text{mL}$), moderate antileishmanial (IC_{50} 13.0 $\mu\text{g}/\text{mL}$) and mild antimalarial activity against *Plasmodium falciparum* (D6 clone) and *P. falciparum* (W2 clone) with IC_{50} values of 2.8 and 2.6 $\mu\text{g}/\text{mL}$, respectively. Compound **5** had moderate antileishmanial activity (IC_{50} 17.0 $\mu\text{g}/\text{mL}$). Compound **9** showed moderate anti-MRSA (IC_{50} 6.5 $\mu\text{g}/\text{mL}$), weak anticandidal (IC_{50} 20.0 $\mu\text{g}/\text{mL}$) and mild antimalarial activity against *P. falciparum* (D6 clone) and *P. falciparum* (W2 clone) with IC_{50} values of 2.8 and 2.0 $\mu\text{g}/\text{mL}$, respectively. Compound **10** exhibited strong antileishmanial activity (IC_{50} 4.5 $\mu\text{g}/\text{mL}$). Compound **11** displayed weak antileishmanial activity (IC_{50} 31.0 $\mu\text{g}/\text{mL}$).

Compounds **2**, **5** and **11** displayed strong, **1**, **7** and **10** moderate and **8** and **9** weak anti-oxidant activities in the DPPH assay, with **6** being inactive.

Compounds **2**, **3** and **4** exhibited no antinociceptive action in both tail-flick and hot-plate assays up to 120 min following injection (Supplementary data).

3. Conclusion

In previous studies (Ahmed et al., 2008; Radwan et al., 2008) we reported the isolation, identification and biological activities of seventeen new cannabinoids from high potency *C. sativa*. In a continuation of this work, the isolation and identification of eleven non-cannabinoid constituents is reported. Among them, **1–6** are new metabolites. Flavones **8** and **9** were identified for the first time from this plant. Compounds **1**, **5**, **10** and **11** displayed antileishmanial activity, while **1** and **9** showed anti-MRSA activity. Compounds **2**, **5**, **7**, **10** and **11** exhibited anti-oxidant activity.

4. Experimental

4.1. General

^1H NMR (400 MHz), ^{13}C NMR (100 MHz), DEPT-135, APT and 2D NMR spectra were recorded using the residual solvent signal as internal standard on a Varian AS 400. IR spectra were measured on a Bruker Tensor 27. UV spectra were obtained on a Varian Cary 50 Bio UV–visible spectrophotometer. Optical rotation was measured on an Autoplot IV automatic polarimeter. High resolution mass spectra were measured using a Bruker BioApex. HPLC was performed on a Waters Delta Prep 4000 Preparative Chromatography System connected to a Waters 486 Tunable UV Absorbance detector using Phenomenex Luna C18 and Si columns (250×21.2 mm, $5 \mu\text{m}$, 100 \AA). GCMS analysis was carried out on a HP 6890 series GC, equipped with a split/splitless capillary injector, a HP 6890 series injector autosampler and an Agilent DB-5 ms column ($30 \text{ m} \times 0.25 \text{ mm} \times 0.25 \mu\text{m}$), interfaced to a HP 5973 mass selective detector (MSD). The injector temperature was 250°C and $1 \mu\text{L}$ injections were performed in the splitless mode, with the splitless time set at 60 s, the split flow set at 50 mL/min and the septum purge valve set to close 60 s after the injection occurred. The oven temperature was raised from 70 to 270°C (hold 20 min) at a rate of $5^\circ\text{C}/\text{min}$, for a total run time of 60 min; the transfer line temperature was 280°C .

4.2. Plant material

C. sativa plants were grown from high potency Mexican seeds. The seeds and plants were authenticated by Dr. Suman Chandra, The University of Mississippi, and a specimen (S1310V1) was deposited at the Coy Waller Complex, The University of Mississippi. Whole buds of mature female plants were harvested, air-dried, packed in barrels and stored at low temperature (-24°C).

4.3. Extraction, isolation and characterization

The plant material (9.0 kg) was sequentially extracted with hexanes (48 L), CH_2Cl_2 (40 L), EtOAc (40 L), EtOH (40 L), EtOH/ H_2O (36 L, 1:1) and H_2O (40 L) at room temperature. The extracts were evaporated under reduced pressure at 40°C to afford hexanes (1.48 kg), CH_2Cl_2 (0.15 kg), EtOAc (0.13 kg), EtOH (0.09 kg), EtOH/ H_2O (0.77 kg) and H_2O (0.54 kg) extracts for a total extract of 3.16 kg (35.1%, w/w). Portions of the CH_2Cl_2 , EtOAc and EtOH extracts were combined (191.0 g) since they showed similar TLC profiles (EtOAc/*n*-hexane, 4:6) and were subjected to silica gel VLC, eluting with EtOAc/*n*-hexane [0:100, 10:90, 20:80, 30:70, 40:60, 50:50, 75:25, 100:0 (2 L of each mixture)] followed by EtOH (4 L), yielding 9 fractions (A–I). Fraction B (10 g) was fractionated on a

silica gel column (EtOAc/petroleum ether, 90:10) to give 16 sub-fractions (B_{1-16}). Subfraction B_1 (265 mg) was purified by Si HPLC (EtOAc/*n*-hexane, 5:95, 25 mL/min, UV 270 nm) to afford **1** (12.3 mg, rt. = 4.0 min). Subfraction B_3 (236.0 mg) was purified by Si-SPE column eluting with MeOH/ CH_2Cl_2 (1:19) yielding **6** (21.6 mg). Fraction D (14.3 g) was subjected to silica CC (EtOAc/petroleum ether, 5:95 to 20:80) followed by C18 flash chromatography (MeOH/ H_2O , 8:2) and C18 HPLC (MeCN/ H_2O , 50:50, 25 mL/min, UV 270 nm) to afford **9** (5.0 mg, rt. = 3.8 min), **10** (1.2 mg, rt. = 4.6 min), **7** (15.0 mg, rt. = 6.7 min) and **2** (11.6 mg, rt. = 8.7 min). Fraction F (28.5 g) was applied to a silica gel column (EtOAc/*n*-hexane, 10:90 to 60:40), yielding 42 fractions (F_{1-42} , 200 mL each). Fraction F_{32-35} (5.5 g) was subjected to silica gel CC (MeOH/ CH_2Cl_2 , 3:97) to yield 24 sub-fractions (SF_{1-24}). SF_{2-3} (210 mg) was applied to a Si-SPE column (EtOAc/*n*-hexane, 10:90) followed by C18 HPLC purification (MeOH/ H_2O , 85:15, 25 mL/min, UV 279 nm) to yield **3** (2.3 mg, rt. = 6.3 min) and **4** (3.9 mg, rt. = 8.8 min). SF_{3-6} (1.9 g) was purified by Sephadex LH-20 CC (MeOH) followed by C18-SPE purification (MeOH/ H_2O , 75:25) to give **5** (12.9 mg), **8** (284.5 mg) and **11** (15.0 mg).

4.3.1. 5-Acetoxy-6-geranyl-3-*n*-pentyl-1,4-benzoquinone (**1**)

Orange amorphous powder; UV λ_{max} (MeOH): 205, 270, 384 nm; IR ν_{max} (neat): 1663 (C=O, ketone), 1780 (C=O, ester), 1610 (C=C) cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ_{H} 0.87 (3H, t, $J = 6.4$ Hz, H_3-5'), 1.32 (4H, m, H_2-3' and H_2-4'), 1.56 (3H, s, H_3-8''), 1.57 (2H, m, H_2-2'), 1.63 (3H, s, H_3-9''), 1.69 (3H, s, H_3-10''), 1.94 (2H, m, H_2-4''), 2.05 (2H, m, H_2-5''), 2.32 (3H, s, OCOCH₃), 2.39 (2H, t, $J = 7.6$ Hz, H_2-1'), 3.10 (2H, d, $J = 7.2$ Hz, H_2-1''), 4.99 (1H, t, $J = 7.2$ Hz, H-2''), 5.02 (1H, t, $J = 7.2$ Hz, H-6''), 6.53 (1H, s, H-2); ^{13}C NMR (CDCl_3 , 100 MHz): δ_{C} 187.1 (C-1), 132.7 (C-2), 148.3 (C-3), 180.8 (C-4), 149.1 (C-5), 135.6 (C-6), 28.9 (C-1'), 27.6 (C-2'), 31.6 (C-3'), 22.6 (C-4'), 14.1 (C-5'), 23.0 (C-1''), 118.4 (C-2''), 138.4 (C-3''), 39.8 (C-4''), 26.7 (C-5''), 124.1 (C-6''), 131.8 (C-7''), 17.9 (C-8''), 25.9 (C-9''), 16.4 (C-10''), 20.5 (OCOCH₃), 168.1 (OCOCH₃); HRESIMS m/z 373.2425 [M+H]⁺ (Calc. for $\text{C}_{23}\text{H}_{33}\text{O}_4$, 373.2379).

4.3.2. 4,5-Dihydroxy-2,3,6-trimethoxy-9,10-dihydrophenanthrene (**2**)

Brown amorphous powder; UV λ_{max} (MeOH): 220, 267, 310 nm; IR ν_{max} (neat): 3420 (OH), 1610, 1537, 1462 (benzene ring) cm^{-1} ; For ^1H and ^{13}C NMR spectroscopic data, see Table 1; HRESIMS m/z 325.1100 [M+Na]⁺ (Calc. for $\text{C}_{17}\text{H}_{18}\text{O}_5\text{Na}$, 325.1052).

4.3.3. 4-Hydroxy-2,3,6,7-tetramethoxy-9,10-dihydrophenanthrene (**3**)

Pale brownish amorphous powder; UV λ_{max} (MeOH): 220, 267, 310 nm; IR ν_{max} (neat): 3420 (OH), 1610, 1537, 1462 cm^{-1} ; For ^1H and ^{13}C NMR spectroscopic data, see Table 1; HRESIMS m/z 339.1279 [M+Na]⁺ (Calc. for $\text{C}_{18}\text{H}_{20}\text{O}_5\text{Na}$, 339.1208).

4.3.4. 4,7-Dimethoxy-1,2,5-trihydroxyphenanthrene (**4**)

Reddish brown amorphous powder; UV λ_{max} (MeOH): 258, 282, 303 nm; IR ν_{max} (neat): 3413 (OH), 1610, 1533, 1462 cm^{-1} ; For ^1H and ^{13}C NMR spectroscopic data, see Table 1; HRESIMS m/z 571.1630 [2M-H]⁻ (Calc. for $\text{C}_{32}\text{H}_{27}\text{O}_{10}$, 571.1604).

4.3.5. 8-Geranyl-5,7,4'-trihydroxy-3'-methoxyflavone (Cannflavin C) (**5**)

Yellow amorphous powder; UV λ_{max} (MeOH): 275, 340, (+NaOMe) 280, 342, 401, (+NaOAc) 280, 340, (+AlCl₃) 300, 346, 360, (+AlCl₃+HCl) 300, 346, 360 nm; IR ν_{max} (neat): 3421 (OH), 1662 (C=O) cm^{-1} ; ^1H NMR (acetone- d_6 , 400 MHz): δ_{H} 1.48 (3H, s, H_3-9''), 1.53 (3H, s, H_3-8''), 1.82 (3H, s, H_3-10''), 1.96 (2H, m, H_2-4''), 2.05 (2H, m, H_2-5''), 3.57 (2H, d, $J = 6.8$ Hz, H_2-1''), 3.98 (3H, s, OMe-3'), 5.02 (1H, t, $J = 6.8$ Hz, H-6''), 5.35 (1H, t, $J = 6.8$ Hz, H-2''), 6.36 (1H, s, H-6), 6.66 (1H, s, H-3), 7.01 (1H, d, $J = 8.0$ Hz, H-5'), 7.58 (1H, d, $J = 2.0$ Hz, H-2'), 7.64 (1H, dd, $J = 2.0, 8.0$ Hz, H-6'),

13.05 (1H, s, HO-5); ^{13}C NMR (acetone- d_6 , 100 MHz): δ_{C} 164.1 (C-2), 103.3 (C-3), 182.7 (C-4), 160.5 (C-5), 98.9 (C-6), 162.4 (C-7), 106.5 (C-8), 155.4 (C-9), 104.5 (C-10), 124.4 (C-1'), 109.7 (C-2'), 148.3 (C-3'), 151.1 (C-4'), 115.8 (C-5'), 120.7 (C-6'), 21.7 (C-1''), 123.0 (C-2''), 135.1 (C-3''), 39.7 (C-4''), 26.7 (C-5''), 124.4 (C-6''), 131.0 (C-7''), 17.9 (C-8''), 25.1 (C-9''), 16.0 (C-10''), 55.8 (OMe-3'); HRESIMS m/z 459.1766 $[\text{M}+\text{Na}]^+$ (Calc. for $\text{C}_{26}\text{H}_{28}\text{O}_6\text{Na}$, 459.1784).

4.3.6. β -Sitosteryl-3-O- β -D-glucopyranoside-2'-O-palmitate (**6**)

White amorphous powder; $[\alpha]_{\text{D}}^{25}$: +91.3 (c 0.05, CHCl_3); IR ν_{max} (neat): 1736 (OH), 1610 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ_{H} 0.63 (9H, m , H₃-26, 27, 29), 0.75 (3H, s , H₃-18), 0.83 (3H, t , $J = 7.2$ Hz, H₃-16''), 0.90 (3H, d , 6.8 Hz, H₃-21), 0.99 (3H, s , H₃-19), 1.24 (18H, bs , H₂-5'' to H₂-13''), 1.28–1.30 (4H, m , H₂-4'', 15''), 2.34 (2H, m , H₂-2''), 3.49 (1H, m , H-3), 3.35–3.82 (4H, H-2' to H-5'), 4.33 (1H, d , $J = 7.6$ Hz, H-1'), 4.1–4.2 (2H, m , H₂-6') 5.33 (1H, bs , H-6); ^{13}C NMR (CDCl_3 , 100 MHz): δ_{C} 37.6 (C-1), 29.6 (C-2), 80.1 (C-3), 39.2 (C-4), 140.6 (C-5), 122.2 (C-6), 32.2 (C-7), 32.1 (C-8), 50.3 (C-9), 36.9 (C-10), 21.3 (C-11), 40.0 (C-12), 42.5 (C-13), 57.0 (C-14), 24.5 (C-15), 28.3 (C-16), 56.4 (C-17), 12.1 (C-18), 19.6 (C-19), 36.4 (C-20), 19.0 (C-21), 34.2 (C-22), 26.4 (C-23), 46.0 (C-24), 29.4 (C-25), 19.2 (C-26), 20.0 (C-27), 23.3 (C-28), 12.2 (C-29), 101.6 (C-1'), 73.5 (C-2'), 76.6 (C-3'), 70.8 (C-4'), 73.8 (C-5'), 64.1 (C-6'), 174.6 (C-1''), 34.5 (C-2''), 25.2 (C-3''), 29.6 (C-4'' to C-13''), 30.0 (C-14''), 22.9 (C-15''), 14.3 (C-16''); HRESIMS m/z 837.6621 $[\text{M}+\text{Na}]^+$ (calc. for $\text{C}_{51}\text{H}_{90}\text{O}_7\text{Na}$, 837.6584).

4.3.7. α -Cannabispiranol (**7**)

Colorless prisms (MeOH/H₂O); mp 182 °C; UV λ_{max} (MeOH): 210, 222 nm; IR ν_{max} (neat): 3410, 3180 (OH), 1610, 1596 cm^{-1} ; ^1H NMR ($\text{C}_5\text{D}_5\text{N}$, 400 MHz): δ_{H} 1.75 (4H, m , H₂-2', 6'), 2.12 (2H, t , $J = 7.2$ Hz, H₂-2), 2.25 (4H, m , H₂-3', 5'), 2.90 (2H, t , $J = 7.2$ Hz, H₂-3), 3.70 (3H, s , OMe-5), 4.10 (1H, m , H-4'), 6.52 (1H, bs , H-4), 6.67 (1H, bs , H-6); ^{13}C NMR ($\text{C}_5\text{D}_5\text{N}$, 100 MHz): δ_{C} 49.2 (C-1), 36.4 (C-2), 31.9 (C-3), 102.0 (C-4), 161.1 (C-5), 101.3 (C-6), 156.7 (C-7), 129.9 (C-8), 146.9 (C-9), 34.6 (C-2', 6'), 34.3 (C-3', 5'), 70.8 (C-4'), 55.6 (OMe-5); HRESIMS m/z 249.2393 $[\text{M}+\text{H}]^+$ (Calc. for $\text{C}_{15}\text{H}_{21}\text{O}_3$, 249.1491); GCMS m/z (rel. int.): 248 (M^+ , 33%), 230 (13%), 215 (12%), 201 (12%), 189 (100%), 176 (65%), 161 (18%).

4.4. Acid hydrolysis of **6**

Compound **6** (5 mg) was refluxed with 3% H₂SO₄ in MeOH (4 mL) for 8 h. The reaction mixture was neutralized with Na₂CO₃ and extracted with EtOAc (3 \times 10 mL). TLC comparison of the aqueous layer with authentic sugar samples (CH₂Cl₂/MeOH/H₂O, 6:4:1; n -BuOH/AcOH/H₂O, 5:5:1) identified the glycone as β -D-glucose.

4.5. Alkaline hydrolysis of **6** and identification of the fatty acid

Compound **6** (5 mg) was added to 10% KOH in MeOH (10 mL) and H₂O (5 mL), and after refluxing for 30 min, H₂O (10 mL) was added, followed by extraction with CHCl₃ (3 \times 10 mL). The aqueous layer was acidified with dil. HCl to pH 5 and extracted with CHCl₃ (3 \times 10 mL). The CHCl₃ layer was dried (Na₂SO₄), filtered and concentrated *in vacuo*. The filtrate was dissolved in dry Et₂O and MeOH (4 drops), followed by methylation with trimethylsilyl-diazomethane (200 μL). The reaction mixture was left open at room temperature for 30 min, concentrated under N₂ and analyzed by GCMS. The fatty acid methyl ester was identified as methyl palmitate via a library search (NIST).

4.6. Antimicrobial, antileishmanial and antimalarial bioassay

The isolated compounds were evaluated for antimicrobial (*Candida albicans* ATCC 90028, *Escherichia coli* ATCC 35218, *Pseudomo-*

nas aeruginosa ATCC 27853, *Mycobacterium intracellulare* ATCC 23068, *Aspergillus fumigat* ATCC 90906, Methicillin Resistant *Staphylococcus aureus* ATCC 43300) (Bharate et al., 2007; Babu et al., 2006), antileishmanial (Radwan et al., 2008) and antimalarial activity [*P. falciparum* (D6 clone) and *P. falciparum* (W2 clone)] (Bharate et al., 2007).

4.7. Anti-oxidant activity

A TLC autographic assay for DPPH (1,1-diphenyl-2-picrylhydrazyl) radical-scavenging effect was used to determine anti-oxidant activity (Takamatsu et al., 2003). The isolated compounds were dissolved in DMF (2 mg/mL) and applied in the form of a spot (4 μL , 4–5 mm in diameter) on silica gel GF plates. The residual DMF was removed under vacuum (15–20 min). A similar amount of vitamin E in DMF was used as positive anti-oxidant control. The radical-scavenging effects of the compounds were detected on the TLC plate using DPPH spray reagent (0.2% w/v in MeOH). The plate was observed 30 min after spraying. Active compounds are observed as yellow spots against a purple background. Relative radical-scavenging activity was assigned as “strong” (compounds that produce an intense bright yellow spot), “medium” (compounds that produce a clear yellow spot), “weak” (compounds that produce a weakly visible yellow spot), or “not active” (compounds that produce no sign of any yellow spot). Vitamin E produced an intense bright yellow spot.

Acknowledgments

The project described was supported in part by Grant No. 5P20RR021929-02 from the National Center for Research Resources and in part by the National Institute on Drug Abuse, Contract # N01DA-5-7746. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center for Research Resources or the National Institutes of Health. We are grateful to Dr. Bharathi Avula for assistance with the HRESIMS, and to Dr. Melissa Jacob, Ms. Marsha Wright, Dr. Babu Tekwani and Dr. Shabana Khan for conducting the antimicrobial and antiprotozoal testing.

Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.phytochem.2008.07.010.

References

- Abegaz, B.M., Ngadjui, B.T., Dongo, E., Tamboue, H., 1998. Prenylated chalcones and flavones from the leaves of *Dorstenia kameruniana*. *Phytochemistry* 49, 1147–1150.
- Agrawal, P.K., 1989. Carbon-13 NMR of Flavonoids, Studies in Organic Chemistry, vol. 39. Elsevier, Amsterdam.
- Ahmed, S.A., Ross, S.A., Slade, D., Radwan, M.M., Zulfikar, F., ElSohly, M.A., 2008. Cannabinoid ester constituents from high-potency *Cannabis sativa*. *J. Nat. Prod.* 71, 536–542.
- Babu, K.S., Li, X.C., Jacob, M.R., Zhang, Q., Khan, S.I., Ferreira, D., Clark, A.M., 2006. Synthesis, antifungal activity, and structure-activity relationships of coruscanone A analogues. *J. Med. Chem.* 49, 7877–7886.
- Bharate, S.B., Khan, S.I., Yunus, N.A.M., Chauthe, S.K., Jacob, M.R., Tekwani, B.L., Khan, I.A., Singh, I.P., 2007. Antiprotozoal and antimicrobial activities of *O*-alkylated and formylated acylphloroglucinols. *Bioorg. Med. Chem.* 15, 87–96.
- Boeren, E.G., ElSohly, M.A., Turner, C.E., Salemink, C.A., 1977. β -Cannabispiranol: a new non-cannabinoid phenol from *Cannabis sativa* L. *Experientia* 33, 848.
- Chang, I.M., Yun, H.S., Yamasaki, K., 1981. Revision of ^{13}C NMR assignments of β -sitosterol and β -sitosteryl-3-O- β -D-glucopyranoside isolated from *Plantago asiatica* seed. *Saengyak Hakhochei* 12, 12–14.
- Choi, Y.H., Hazekamp, A., Peltenburg-Looman, A.M.G., Frederich, M., Erkelens, C., Lefebvre, A.W.M., Verpoorte, R., 2004. NMR assignments of the major

- cannabinoids and cannabiflavonoids isolated from flowers of *Cannabis sativa*. *Phytochem. Anal.* 15, 345–354.
- Crombie, L., Crombie, W.M.L., Jamieson, S.V., 1979. Isolation of cannabispiradienone and cannabidiolphenanthrene biosynthetic relationships between the spirans and dihydrostilbenes of Thailand cannabis. *Tetrahedron Lett.* 20, 661–664.
- Crombie, L., Crombie, W.M.L., 1982. Natural products of Thailand high Δ^1 -THC-strain *Cannabis* the bibenzyl-spiran-dihydrophenanthrene group: relations with cannabinoids and canniflavones. *J. Chem. Soc., Perkin Trans. 1*, 1455–1466.
- El-Feraly, F.S., El-Sherei, M.M., Al-Muhtadi, F.J., 1986. Spiro-indans from *Cannabis sativa*. *Phytochemistry* 25, 1992–1994.
- ElSohly, M.A., Slade, D., 2005. Chemical constituents of marijuana: the complex mixture of natural cannabinoids. *Life Science* 78, 539–548.
- Ishii, H., Seo, S., Tori, K., Tozoy, T., Yoshimura, Y., 1977. The structures of saikosaponin-E and acetylsaikosaponins, minor components isolated from *Bupleurum falcatum* L. determined by ^{13}C NMR spectroscopy. *Tetrahedron Lett.* 18, 1227–1230.
- Kovganko, N.V., Nashkan, Z.N., Borisov, E.V., 2000. ^{13}C NMR spectra of functionally substituted 3β -chloro derivatives of cholesterol and β -sitosterol. *Chem. Nat. Compd.* 36, 595–598.
- Leong, Y.W., Kang, C.C., Harrison, L.J., Powell, A.D., 1997. Phenanthrenes, dihydrophenanthrenes and bibenzyls from the orchid *Bulbophyllum vaginatum*. *Phytochemistry* 44, 157–165.
- Mabry, T.J., Markham, K.R., Thomas, M.B., 1970. *The Systematic Identification of Flavonoids*. Springer-Verlag, New York.
- Mossa, J.S., Muhammad, I., Ramadan, A.F., Mirza, H.H., El-Feraly, F.S., Hufford, C.D., 1999. Alkylated benzoquinone derivatives from *Maesa lanceolata*. *Phytochemistry* 50, 1063–1068.
- Radwan, M.M., Ross, S.A., Ahmed, S.A., Slade, D., Zulfiqar, F., ElSohly, M.A., 2008. Isolation and characterization of new cannabis constituents from a high-potency variety. *Planta Med.* 74, 267–272.
- Ross, S.A., ElSohly, M.A., 1995. Constituents of *Cannabis sativa* L XXVII. A review of the natural constituents: 1980–1994. *Zagazig J. Pharm. Sci.* 4, 1–10.
- Réthy, B., Kovács, A., Zupkó, I., Forgo, P., Vasas, A., Falkay, G., Hohmann, J., 2006. Cytotoxic phenathrenes from the rhizome of *Tamus communis*. *Planta Med.* 72, 767–770.
- Segre, A., Mannina, L., 1997. ^1H NMR study of edible oils. *Recent Res. Develop. Oil Chem.* 1, 297–308.
- Shoyama, Y., Nishioka, I., 1978. Cannabis. XIII. Two new spiro-compounds, cannabispinol and acetyl cannabispinol. *Chem. Pharm. Bull.* 26, 3641–3646.
- Stermitz, F.R., Suess, T.R., Schauer, C.K., Anderson, O.P., Bye, R.A., 1983. New and old phenanthrene derivatives from *Oncidium cebolleta*, a peyote-replacement plant. *J. Nat. Prod.* 46, 417–423.
- Takamatsu, S., Hodges, W.T., Rajbhandari, I., Gerwick, W.H., Hamann, M.T., Nagle, D.G., 2003. Marine natural products as novel anti-oxidant prototypes. *J. Nat. Prod.* 66, 605–608.
- Takemoto, T., Hikino, Y., Nomoto, K., Hikino, H., 1967. Structure of cyasterone, a novel C29 insect-moulting substance from *Cyathula capitata*. *Tetrahedron Lett.* 33, 3191–3194.
- Terui, Y., Tori, K., Tsuji, N., 1976. Esterification shifts in carbon-13 NMR spectra of alcohols. *Tetrahedron Lett.* 8, 621–622.
- Toth, L., Bulyaki, M., Bujtas, G., 1980. Flavonoids from *Odontites rubra* (Baumg.). *Opiz. Pharmazie.* 35, 335–336.
- Turner, C.E., ElSohly, M.A., Boeren, E.G., 1980. Constituents of *Cannabis sativa* L XVII. A review of the natural constituents. *J. Nat. Prod.* 43, 169–234.
- Yamasaki, K., Kasai, R., Masaki, Y., Okihara, M., Tanaka, O., Oshio, H., Takagi, S., Yamaki, M., Masuda, K., Nonaka, G., Tsuboi, M., Nishioka, I., 1977. Application of ^{13}C NMR to the structural elucidation of acylated plant glycosides. *Tetrahedron Lett.* 14, 1231–1234.